
Adolescent Health

Prescription Pain Medicine Misuse





Data Source: Alaska Division of Public Health, Alaska Youth Risk Behavior Surveillance System Table C-95

Definition

Among high school students (grades 9-12) prescription pain medicine misuse is defined as having used a prescription pain medication (such as OxyContin, Percocet, Vicodin, codeine, or Hydrocodone) without a doctor's prescription or differently than prescribed one or more times during their life.

Summary

- » In 2019, 13.7% of Alaska Native high school students reported having used prescription pain medications without a doctor's prescription or differently than prescribed in their lifetime.
- In 2019, there was no statistically significant difference between Alaska Native and Alaska White high school students in prescription pain medicine misuse.

Adolescent Health





Table C-95: High School Student Prescription Pain Medicine Misuse, 2017-2019

Data Source: Alaska Division of Public Health, Alaska Youth Risk Behavior Surveillance System

| | | Alaska Native Students Statewide | Alaska White Students Statewide |
|------|---------------------|-------------------------------------|------------------------------------|
| 2017 | n | 343 | 569 |
| | % | 16.2% | 15.2% |
| | Confidence Interval | (11.9%-21.7%) | (12.1%-19.0%) |
| 2019 | n | 425 | 903 |
| | % | 13.7% | 14.0% |
| | Confidence Interval | (9.5%-19.3%) | (11.1%-17.5%) |